

Bacteriological Profile and Antimicrobial Susceptibility Pattern of Isolates from Ocular Infections among the Patients Attending Ophthalmology Department of a Tertiary Care Hospital

Indira Ananthapadmanabhasamy^{1*}, Shambhavi GK², CH Srinivasa Rao¹

INTRODUCTION

Bacteria are the major cause of ocular infections around the world. Infection may either be due to single microbe or polymicrobial and is mostly associated with factors like age, contact lens usage, trauma, surgery, poor ocular hygiene or any prior ocular pathology [1]. Ocular infections manifest commonly in the form of blepharitis, conjunctivitis, keratitis, endophthalmitis, orbital cellulitis or dacryocystitis [2]. The source of eye infection may be exogenous which is the most common mode or endogenous invasion of the micro-organisms which can be carried in the bloodstream. External or exogenous bacterial infection of the eye are usually localised to some extent or may spread to the adjacent tissues [3].

The eye is normally protected by the continuous flow of tears and the blink reflex which mechanically prevents foreign substances invading it. Tear film also possess few enzymes like lysozyme, secretory IgA, lactate dehydrogenase, beta-lysin, defensins, etc which are present at high levels in tears that is capable of reducing microbial colonisation on the ocular surface thereby preventing bacterial infections [4]. The virulence factors of pathogenic microorganisms and impaired host resistance favours the ocular infections. The common organisms that cause eye infections are Propriobacterium acne, Staphylococcus aureus, Streptococcus pneumoniae, Moraxella spp [2]. Conjunctivitis, inflammation of the mucosa of conjunctiva, is the most frequent ocular case with economic and social burdens [5]. Blepharitis which is an inflammation of the eyelid can cause loss of eye lash [6]. Keratitis, the most serious eye infection is the leading cause of corneal blindness. Moreover, the disease can also progress to endophthalmitis if not diagnosed early [7]. Dacryocystitis is an inflammation of the nasolacrimal duct [8]. Exogenous endophthalmitis is an infective complication of primary cataract, intraocular surgery and ocular trauma due to the introduction of infectious pathogens like bacteria whereas the endogenous one is commonly due to systemic dissemination of the pathogens [9].

The eye infections if left untreated can damage the structures of the eye, causing visual impairment and blindness [1]. Knowledge on the specific aetiology is essential for the effective management of ocular infections. However these infections are mainly managed empirically and very less is known about the specific aetiology [10, 11]. The purpose of this study is to identify the specific bacterial pathogen responsible for the development of particular ocular infection and to determine their in-vitro susceptibility to commonly used antibacterial agents in clinical practice.

METHODOLOGY

A hospital based cross sectional study was conducted among the patients attending the ophthalmology department in a tertiary care hospital, Tirupati from October 2019 to September 2020 with the approval from the Institutional Ethical Committee. All the patients attending the Ophthalmology Department with ocular infections were included in the

study. Patients who were already on antibacterial therapy were excluded from the study.

OPERATIONAL DEFINITIONS

Conjunctivitis: Inflammation or infection of the conjunctiva [12]

Blepharitis: Inflammation or infection of the eyelid margin [13]

Dacryocystitis: Inflammation of the lacrimal sac [13]

Infective uveitis: Inflammation or infection of the uveal tissue of the eye [13]

Endophthalmitis: Endophthalmitis is a purulent inflammation of the intraocular fluids [vitreous and aqueous] usually due to infection [13]

DATA COLLECTION

The study included 290 patients with ocular infections who attended the Ophthalmology Department of tertiary care hospital, Tirupati. The socio-demographic data of each study participants were collected. The ocular examination was performed using slit lamp bio-microscopy to identify any focus of infection and inflammation, to come to a provisional diagnosis. The provisional diagnosis was recorded and the specimen was collected by the attending ophthalmologist using a standard protocol from the study participants.

SPECIMEN COLLECTION

Conjunctival swab: Swabs are collected prior to the start of anti-microbial therapy. After the clinical examination with the help of a sterile cotton swab the discharge collected in the conjunctival cul-de-sac and lacus lacrimalis is collected by a gentle stroke from lateral canthus to medial canthus. This procedure is done without the use of any anaesthetic drops. Two such swabs are collected from each eye and sent for microbiological examination [13]

Corneal scrapping: Two drops of 0.5 % proparacaine is instilled in the lower fornix of the affected eye. A lid speculum is applied to separate the eyelids. Under slit lamp or operating microscope the affected eye is examined using direct illumination and any debris or mucus is cleaned using a sterile swab. Kimura spatula or a number 15 Bard Parker knife is used to scrape the leading edge and base of the ulcer. The specimen was directly inoculated into the blood agar plate. Multiple specimens were obtained to enhance the yield [14]. Smears are prepared by transferring the specimen on to the glass slide over an area of approximately 1 cm in diameter for microscopic examination.

Anterior chamber paracentesis: After a local anaesthetic drops of 0.5% proparacaine instilled in lower fornix of the affected eye, a 26 or 30 gauge needle with insulin or 2 ml syringe was used for sample collection. Lid speculum was used to separate the eyelids. Under operating microscope or slit lamp needle was entered in to the anterior chamber by a valvular self-sealing paracentesis obliquely through the stroma via the lower limbus. 0.1

¹Department of Microbiology, Sri Venkateswara Medical College, Tirupati, India

²Department of Ophthalmology, Sri Venkateswara Medical College, Tirupati, India

*Correspondence to: Indira Ananthapadmanabhasamy, Department of Microbiology, Sri Venkateswara Medical College, Tirupati, India, Tel: 8374149975; Email: drindhustar@gmail.com

Received date: May 27, 2021; Accepted date: September 15, 2021; Published date: September 25, 2021



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact reprints@pulsus.com

to 0.3 ml of aqueous was collected avoiding damage to endothelium or the lens. The needle was withdrawn and an external pressure was applied to the entrance with sterile cotton tip applicator. A drop of antibiotic was instilled in the conjunctival sac and eye was patched [15].

Specimen/swabs collected were labelled and transported immediately to microbiology laboratory

CULTURE AND IDENTIFICATION

The first swab was subjected to Gram stain where the presence of bacteria its Gram’s reaction and presence of pus cells were assessed.

The second swab was inoculated on to 5% sheep blood agar, MacConkey agar, and Mannitol salt agar and inoculated aerobically at 37°c for 24 hours.

All the plates were examined for pure growth. The pure colonies were obtained for further identification methods using standard microbiological techniques, i.e Gram stain, colony morphology and biochemical tests.

The identification of Gram positive bacteria was based on the type of haemolysis on sheep blood agar followed by catalase test, coagulase test, bile solubility and Optochin disk test [16, 17].

The identification of Gram negative bacteria was based on catalase test, oxidase test, hanging drop motility test, indole test, Methyl red test, Voges Proskauer test, Citrate utilisation test, triple sugar iron agar test, urease test and Arginine ornithine decarboxylase test[18].

Antimicrobial susceptibility testing [AST]

AST was carried out for each bacterial isolate using Kirby Bauer’s disc diffusion technique based on CRSI [Clinical and Laboratory Standard Institute] guidelines [19]

Briefly, 3 to 5 pure colonies of bacteria were transferred into a test tube containing 1 ml of sterile normal saline mixed homogenously and adjusted to 0.5 Mc Farland standards.

The suspension was inoculated on to Mueller Hilton agar [MHA] for non-fastidious organism and in 5% sheep blood agar for fastidious like Streptococcus pneumoniae.

Each antibiotic disc was placed manually on agar plate and incubated at 37 °c for 24 hours and the zone of inhibition around the disc was measured to the nearest milli-meter using graduated scale. The isolate were classified as susceptible, intermediate and resistance according to CLSI guidelines.

As there are no antibiotic susceptibility break points for topical antibiotic therapy higher antibiotic concentration comparatively are achieved in ocular tissue while using topical therapy.

RESULTS

In the present study a total of 290 patients had attended ophthalmology department for ocular infections. Among the study participants 164 [56.52%] were males, and 174 [60%] belongs to rural areas. The most affected age group belongs to > 60 [40.41%] years.

TABLE 1: GENDER WISE DISTRIBUTION OF THE STUDY PARTICIPANTS

Gender	Males	Females	Total
Total No. Of Study Participants	164 [56.55%]	126 [43.45%]	290

CHART 1: RESIDENCE WISE DISTRIBUTION OF THE STUDY PARTICIPANTS




TABLE 2: AGE WISE DISTRIBUTION OF THE STUDY PARTICIPANTS

Age Group	Total Participants [%]	Positive For Bacterial Culture [%]
20-Jan	37 [12.76]	14 [9.54]
21-40	56 [19.31]	28 [19.18]
41-60	83 [28.12]	45 [30.28]
>60	114 [39.31]	59 [40.41]
TOTAL	290	146

CLINICAL DATA

The various clinical conditions seen among 290 patients were conjunctivitis 108 [37.24%], keratitis 37 [12.76%] dacryocystitis 53 [18.28%], blepharitis 52 [17.93%], trauma 24 [8.28%], infective uveitis 10 [3.45%] and endophthalmitis 6 [2.07%].

TABLE 3: DISTRIBUTION OF CLINICAL CONDITIONS AMONG STUDY PARTICIPANTS.

Types Of Presentations	Clinical Presentation	Clinical Presentation	Frequency Of Bacteria Among Clinical Presentation
		n [%]	n [%]
Conjunctivitis		108 [37.24%]	79 [73.14%]
Corneal Ulcer		37 [12.76%]	11 [29.73%]
Dacryocystitis		53 [18.28%]	21 [39.62%]
Periocular +Trauma	Burns	24 [8.28%]	6 [25%]
Infective Uveitis		10 [3.45]	-
Endophthalmitis		6 [2.07%]	1 [16%]
Blepharitis		52 [17.93%]	28 [53.85%]
TOTAL		290	146

BACTERIAL AETIOLOGY OF OCULAR INFECTION

Among the 290 participants with ocular infections, 146 [50.34%] were positive for culture. Among the total isolated, 110 [75.34%] were Gram positive and 36 [24.66%] were Gram negative. The predominant bacteria among the isolate was Staphylococcus aureus [Table 4].

TABLE 4: DISTRIBUTION OF BACTERIA ISOLATED FROM STUDY SPECIMENS

Bacteria	Number [n=146]	Percentage
----------	----------------	------------

Staphylococcus aureus	62	42.47
Coagulase negative Staphylococcus	35	23.97
Streptococcus pneumoniae	13	8.9
Escherichia coli	11	7.53
Klebsiella pneumoniae	13	8.9
Pseudomonas aeruginosa	4	2.74
Moroxella spp	5	3.42
Acinetobacter spp	3	2.05

The most common organism isolated among blepharitis specimens was coagulase negative Staphylococcus and in other clinical conditions Staphylococcus aureus was the most common organism to be isolated [Table 5].

TABLE 5: FREQUENCY OF BACTERIA ISOLATED AMONG THE VARIOUS CLINICAL CONDITIONS

Clinical Condition	BACTERIA ISOLATED							
	S.aureus	CONS	S.pneumoniae	E.coli	K.pneumoniae	P.aeruginosa	Moroxella spp	Acinetobacter spp
Conjunctivitis	34	23	6	5	7	1	3	-
Corneal Ulcer	5	3	-	-	-	2	-	1
Dacryocystitis	17	-	-	2	1	-	1	-
Trauma	-	-	2	1	1	1	-	1
Endophthalmitis	-	-	-	-	-	-	-	1
Blepharitis	6	9	5	3	4	-	1	-
TOTAL	62	35	13	11	13	4	5	3

ANTIMICROBIAL SUSCEPTIBILITY PROFILE

Among the 110 Gram positive bacteria isolated, the most sensitive antibiotic was ciprofloxacin [86.36%] followed by gentamycin [82.72%] and clindamycin [80%] [Table 6]

TABLE 6: ANTIMICROBIAL SUSCEPTIBILITY PATTERN OF THE GRAM POSITIVE BACTERIA ISOLATED

BACTERIA [n=110]	AMX	COT	E	CL	NFX	CIP	TE	GE	VA
S.aureus [n=62]	8	43	13	57	5	53	11	47	62
CONS [n=35]	3	21	17	31	12	29	4	31	35
S.pneumoniae	13	1	13		13	13	13	13	-

[n=13]

Total [n=110]	24	65	43	88	30	95	28	91	97
	-21.8	-59.0	-39.0	-80	-27.2	-86.3	-25.4	-82.7	-10
	2	9	9	%	7	5	5	2	0

AMX: Amoxicillin, CONS: Coagulase Negative Staphylococcus, CIP: Ciprofloxacin, CL: Clindamycin, COT: Cotrimoxazole, E: Erythromycin, GE: Gentamycin, NFX: Norfloxacin, TE: Tetracycline, VA: Vancomycin

Among the 62 Staphylococcus aureus isolated the highest resistance was seen with amoxicillin 54 [87.09%] followed by tetracycline 51 [82.26%] and erythromycin 49 [79.03%].

The total number of Streptococcus pneumoniae isolated were 100% sensitive to amoxicillin, erythromycin and gentamycin.

Among the 36 Gram negative bacteria isolated, 35 [97.22%], 33 [91.67%], and 31 [86.11%] were sensitive to gentamycin, ciprofloxacin, ceftriaxone respectively. [Table 7]

The Gram negative bacteria showed high resistance to tetracycline 7[19.44%], followed by amoxicillin clavulanic acid [30.56%].

TABLE 7: ANTIMICROBIAL SUSCEPTIBILITY PATTERN OF THE GRAM NEGATIVE BACTERIA ISOLATED

BACTERIA	AMC	COT	CIP	CTR	GE	TE	NFX
Escherichia coli [n=11]	9	7	11	9	11	3	6
Klebsiella pneumoniae [n=13]	7	8	13	11	13	1	7
Pseudomonas aeruginosa [n=4]	1	-	2	4	4	-	1
Acinetobacter [n=3]	-	-	3	2	3	-	1
Moroxella [n=5]	1	2	4	5	4	3	1
Total [n=36]	11	17	33	31	35	7	16
	-30.56	-47.22	-91.67	-86.11	-97.22	-19.44	-44.44

AMC: Amoxicillin - Clavulanic acid, CIP: Ciprofloxacin, COT: Cotrimoxazole, CTR: Ceftriaxone, GE: Gentamycin, NFX: Norfloxacin, TE: Tetracycline

DISCUSSION

The present study includes 290 specimens, among which 146 [50.34%] specimens were culture positive with coincides with various study conducted by Mohammed et al [20] [59.6%], Bharathi et al [21] [58.8%] and Shiferaw et al [22] [59.4%].

In the present study the men were more susceptible for ocular infections [56.55%], which is similar to the study conducted by Mohammad et al [20] [53.3%] and Nuzhat et al [23] [54%].

The most common age group affected by ocular infections in the present study is > 60 years [40.41%] which correlates with the study by Shiferaw et al [22] [44.4%].

The ocular infections are more prevalent in rural areas [60%] in the present study which correlates with other studies namely Mohammed et al [20] [67.2%], Shiferaw et al [22] [75%].

In the present study the most common ocular infection reported was conjunctivitis [37.24%], which is similar to other studies by Mohammed et al [20] [32.8%], Shiferaw et al [23] [43.1%], Nuzhat et al [23] [42%] and Hemavathi et al [16] [52%].

The most common organism isolated from the specimens are *Staphylococcus aureus* [42.4%] followed by Coagulase negative *Staphylococcus* [23.97%].

These findings correlate with many other study conducted by Mohammed et al [20] [37.4%, 28.8%], Namitha et al [24] [32.8%, 25%], Nuzhat et al [23] [36%, 20%] by *Staphylococcus aureus* and Coagulase negative *Staphylococcus* respectively.

In the present study, the Gram negative bacteria most commonly reported is *K.pneumoniae* [8.90%] which correlates with the study performed by Namitha et al [24] [6.2%], Muluze et al [25] [14.5%].

Among the *Staphylococcus aureus* [62] isolated in the present study, methicillin sensitive *Staphylococcus aureus* strains were 12.90% while methicillin resistance *Staphylococcus aureus* were 87.10% and all MRSA strains were 100% sensitive to vancomycin.

Among the coagulase negative *Staphylococcus* [35], only 3 isolates showed sensitivity to methicillin, while rest 32 isolates were resistant to it. These isolates were sensitive to vancomycin [100%]. These correlate with the most of the study around the world i.e Shiferaw et al [22], Nuzhat et al [23], Bharathi et al [21].

The *Streptococcus pneumoniae* [13] isolates in the present study were 100% sensitive to amoxicillin, ciprofloxacin, erythromycin, gentamycin which is similar to Mohammad et al [20], Hemavathi et al [16].

Among the Gram negative bacteria [36%] isolated in the present study, the highest sensitivity was towards gentamycin [9.22%] followed by ciprofloxacin [91.67%] which coincides with the study by Mohammed et al [20]

Among the Gram Negative bacterial isolates, *Klebsiella pneumoniae* [13], *Pseudomonas aeruginosa* [4], *Moraxella* [5], showed 100% sensitive to ceftriaxone which is similar to studies by Mohammed et al [20]

In the present study the highest resistance among Gram positive and Gram negative bacteria was shown for amoxicillin [21.82%] and tetracycline [19.44%] respectively.

This is similar to the study performed by Shiferaw et al [22], Mohammed et al [20], and Muluze et al [25].

CONCLUSION

The most common ocular infection reported in our hospital was conjunctivitis followed by dacryocystitis. *Staphylococcus aureus* stands as the most common bacteria to cause ocular infection in the community. Most of the organisms are sensitive to ciprofloxacin and gentamycin and least sensitivity to amoxicillin and tetracycline. In order to prevent the increasing rate of Antimicrobial resistance, identification of bacteria followed by antimicrobial sensitivity testing should be made mandatory in routine practice.

REFERENCES

1. Teweldemedhin M, Gebreyesus H, Atsbaha AH, Asgedom SW, Saravanan M. Bacterial profile of ocular infections: a systematic review. *BMC Ophthalmol* [Internet]. 2017 Nov 25 [cited 2021 May 8]; 17. Available from:
2. Bertino JS. Impact of antibiotic resistance in the management of ocular infections: the role of current and future antibiotics. *Clin Ophthalmol*. 2009;3:507-21.
3. Ramesh S, Ramakrishnan R, Bharathi MJ, Amuthan M, Viswanathan S. Prevalence of bacterial pathogens causing ocular infections in South

- India. *Indian Journal of Pathology and Microbiology*. 2010 Apr 1;53(2):281.
4. Khosravi AD, Mehdinejad M, Heidari M. Bacteriological findings in patients with ocular infection and antibiotic susceptibility patterns of isolated pathogens. *Singapore Med J*. 2007 Aug;48(8):741-3.
5. Cervantes LJ, Mah FS. Clinical use of gatifloxacin ophthalmic solution for treatment of bacterial conjunctivitis. *Clin Ophthalmol*. 2011;5:495-502.
6. Kowalski RP, Dhaliwal DK. Ocular bacterial infections: current and future treatment options. *Expert Rev Anti Infect Ther*. 2005 Feb;3(1): 131-9.
7. Cao J, Yang Y, Yang W, Wu R, Xiao X, Yuan J, et al. Prevalence of infectious keratitis in Central China. *BMC Ophthalmology*. 2014 Apr 2;14(1):43.
8. Maheshwari R, Maheshwari S, Shah T. Acute dacryocystitis causing orbital cellulitis and abscess. *Orbit*. 2009;28(2-3):196-9.
9. Vaziri K, Schwartz SG, Kishor K, Flynn HW. Endophthalmitis: state of the art. *Clin Ophthalmol*. 2015 Jan 8;9:95-108.
10. Baron EJ, Miller JM, Weinstein MP, Richter SS, Gilligan PH, Thomson RB, et al. A guide to utilization of the microbiology laboratory for diagnosis of infectious diseases: 2013 recommendations by the Infectious Diseases Society of America (IDSA) and the American Society for Microbiology (ASM)(a). *Clin Infect Dis*. 2013 Aug;57(4):e22-121.
11. Harbarth S, Samore MH. Antimicrobial Resistance Determinants and Future Control. *Emerg Infect Dis*. 2005 Jun;11(6):794-801.
12. Parson's "Diseases of the Eye." *Br J Ophthalmol*. 1954 Mar;38(3):190.
13. American Academy of Ophthalmology Launches EyeWiki [Internet]. American Academy of Ophthalmology. 2010 [cited 2021 May 27]. Available from:
14. Sharma N, Vajpayee RB, Taylor HR, Laibson PR. Corneal ulcers: diagnosis and management. New Delhi, India: Jaypee Brothers Medical Publishers; 2008.
15. Hemavathi, Sarmah P, Shenoy P. Profile of Microbial Isolates in Ophthalmic Infections and Antibiotic Susceptibility of the Bacterial Isolates: A Study in an Eye Care Hospital, Bangalore. *J Clin Diagn Res*. 2014 Jan;8(1):23-5.
16. Hesje CK, Sanfilippo CM, Haas W, Morris TW. Molecular epidemiology of methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* isolated from the eye. *Curr Eye Res*. 2011 Feb; 36(2):94-102.
17. Collee JG, Duguid JP, Fraser AG, Marmion BP, editors. Mackie and McCartney Practical Medical Microbiology. 13th ed. Vol. 2. London: Churchill Livingstone; 1989.
18. Clinical & Laboratory Standards Institute: CLSI Guidelines [Internet]. Clinical & Laboratory Standards Institute. [cited 2021 May 27]
19. Mohammed AA, Ali MM, Zenebe MH. Bacterial etiology of ocular and periocular infections, antimicrobial susceptibility profile and associated factors among patients attending eye unit of Shashemene comprehensive specialized hospital, Shashemene, Ethiopia. *BMC Ophthalmol*. 2020 Dec;20(1):124.
20. Bharathi MJ, Ramakrishnan R, Shivakumar C, Meenakshi R, Lionraj D. Etiology and antibacterial susceptibility pattern of community-acquired bacterial ocular infections in a tertiary eye care hospital in south India. *Indian J Ophthalmol*. 2010;58(6):497-507.
21. Shiferaw B, Gelaw B, Assefa A, Assefa Y, Addis Z. Bacterial isolates and their antimicrobial susceptibility pattern among patients with external ocular infections at Borumeda hospital, Northeast Ethiopia. *BMC Ophthalmol*. 2015 Dec;15(1):103.
22. Nuzhat Q, Bandyopadhyay A, Chaurasia R, Singh S, Singh M. Analysis of antimicrobial susceptibility pattern of ocular infections at Regional Ophthalmic Institute in India. *International Journal of Basic & Clinical Pharmacology*. 2020 Mar 24;9:642.
23. Department Of Microbiology, Navodaya Medical College & Reserch Centre, Raichur, B.N N, kshmi M, Rao A. Aerobic Bacteriological

Profile in Cases of Ocular Infections in a Tertiary Care Hospital (Navodaya Medical College & Research Centre, Raichur). IOSRJDMS. 2014;13(11):14-21.

samples at Gondar University Hospital, Northwest Ethiopia. BMC Research Notes. 2014 May 12;7(1):292.

24. Muluye D, Wondimeneh Y, Moges F, Nega T, Ferede G. Types and drug susceptibility patterns of bacterial isolates from eye discharge